# The BASES Expert Statement on the Effectiveness of Vision Training **Programmes**

Produced on behalf of the British Association of Sport and Exercise Sciences by Dr Nicholas Smeeton, Dr Jenny Page, Dr Joe Causer, Dr Mark Wilson, Dr Rob Gray and Prof Mark Williams.

#### Introduction

Two types of vision training programmes exist, namely, generalised vision training (GVT) and sport-specific vision training (SVT). GVT programmes are designed to improve basic visual function (e.g., depth perception, motion perception and peripheral vision). A range of exercises is typically used by vision specialists such as optometrists and ophthalmologists. Although these specialists usually work to assist individuals with visual deficiencies, more recently the same methods have been used with athletes in an effort to improve sports performance. Whilst there is anecdotal support for the use of GVT programmes, there remains a paucity of empirical evidence to suggest that such training improves sports performance. Conversely, research on SVT has been shown to lead to task-specific improvements in sports performance (Smeeton et al., 2005; Williams et al., 2002). Here, we summarise scientific research that examines the utility of both types of training programmes for performance enhancement in sport.

#### Vision training

The rationale for using GVT is that improving basic aspects of visual function will lead to improvements in the performance of perceptual-motor skills that use the functions being trained. For example, exercises that improve an individual's general ability to detect objects in peripheral vision will improve skills such as playing basketball. In GVT it is not necessary to include/represent the specific sport context as part of the training intervention (i.e., sportspecific images are typically not employed).

In contrast, SVT aims to improve the ability to detect, discriminate, and/or identify the specific sources of visual information involved in a particular athlete's sport. For example, rather than using abstract stimuli to train an individual's ability to detect the direction of a moving object, sport-specific stimuli are employed to improve a goalkeeper's ability to detect the direction of motion of a shooter's hip during a football penalty kick.

The first stage in the development of SVT programmes typically involves using experts from within the sport to identify the specific sources of information used during successful performance on the task (i.e., the expert-model approach). Next, interventions are developed that use sport- and task-specific video simulations to ensure that these specific information sources are captured. In situ instructions are given to highlight the most useful places to extract information from the visual display and link this information to actions and outcomes. The effectiveness of these training interventions relies on the development of a comparable knowledge base and visual strategy to that employed by the expert model(s).

### Support for generalised vision training

The evidence used to support GVT is limited due to the fact that very few researchers have demonstrated a causal link between improvements in visual function and subsequent changes in sports

performance. Indeed, the majority of researchers have employed correlational designs. In the USA, Clark et al. (2012) examined whether GVT exercises improved the batting performance of baseball players over two consecutive seasons in the National College Athletic Association. The GVT programme included exercises designed to improve depth perception, saccadic eye movements, accommodation and vergence. Significant increases were reported for several performance metrics following training. While such findings suggest a possible link between GVT and improvements in sports performance, the absence of adequate control and placebo groups precludes definitive conclusions from

The majority of intervention studies using GVT programmes in the sports domain do not support the utility of GVT (see Abernethy, 1996; Williams & Grant, 1999 for reviews). Schwab and Memmert (2012) reported that a group of young field hockey players who participated in a 6-week intervention that included practice using a Dynavision D2® Trainer, Eyeport, Vision Performance Enhancement Program, Hart Charts and P-Rotator improved performance on the same visual test on which they trained. However, there was no improvement on a functional field of view task or an additional measure of transfer (multiple object tracking). Using a randomised, placebo-controlled design, Abernethy and Wood (2001) reported that while participants who underwent one of two GVT programmes did improve performance in a stationary sport-specific transfer test (by 7.25 %) participants in a control group who received no vision training also improved (by 3.3 %). To provide stronger support for the efficacy of GVT programmes, researchers need to ensure that appropriate control and experimental groups are employed so that a cause and effect relationship between GVT and performance can be ascertained.

While there is a lack of evidence for the efficacy of GVT programmes in improving sports performance, such interventions may be useful to redress imbalances (e.g., eye dominance) or deficits in normal visual functioning. The tests used in many GVT programmes could be valuable for screening/testing vision in sport. There may be instances when the visual system of an athlete is not functioning properly and, as a result, sports performance (and general health) might suffer (e.g., Goodrich et al., 2013). GVT for screening and health purposes must not be ignored.

## Support for sport-specific vision training

When sport-specific visual stimuli are used there is some evidence that training does improve sports performance (see Causer et al., 2012 for a review). Williams et al. (2002) trained anticipation of tennis groundstrokes using film-based sport-specific vision (perceptual) training. They demonstrated that anticipation could be trained through video feedback of key visual stimuli from the opponent's action. Not only did anticipation performance improve above that seen in a matched-ability intervention group, but these improvements transferred to an on-court test of anticipation

(the training group's mean responses were 0.187 s quicker than reported for the control and placebo groups). Hopwood et al. (2011) demonstrated that highly skilled cricket players who received visual-perceptual training in conjunction with on-field training, demonstrated greater improvements in in situ fielding tests (catching success improved by 21.7 % from preto post-test) compared to those who received on-field training alone (catching success improved by 16.2 % from pre- to post-test).

Additionally, studies examining eye-gaze

behaviour in sports requiring accurate aiming have shown that skilled performance is linked to having a longer and earlier 'quiet eye' (final fixation prior to the critical movement) on the relevant target location. Training interventions designed to increase this quiet eye dwell time not only successfully enhance task performance (above technically-focused interventions) in laboratory settings, but also transfer to competitive sports settings (see Causer et al., 2012; Vine et al., 2012 for reviews). An important advantage of quiet eye training is that the eye movements are trained in situ. However, as with other SVT programmes, the performance improvements routinely do not generalise to other tasks (Smeeton et al., 2005; Williams et al., 2002). In conjunction with the lack of support for GVT, these findings suggest that benefits arise due to specific ('software') rather than generalised ('hardware') improvements (Abernethy & Wood, 2001). Without task-specific knowledge about the salient visual information, the benefits of having more effective general visual functioning in individuals with already healthy visual function are unlikely to be realised.

#### **Conclusions and recommendations**

- · Generalised vision testing is useful in screening for deficits in visual functioning. Such tests should be conducted by qualified practitioners (e.g., optometrists, ophthalmologists).
- While GVT may improve an athlete's performance on a general test of visual function, there is no peer-reviewed evidence to support the transfer of this improvement to sports performance.
- SVT has been shown to have performance advantages when compared to control and placebo groups across a range of sports. These advantages appear to be task-specific.
- When considering evidence for any vision training intervention, it is important to gauge whether good practice has been followed. Have placebo and control groups been used and has transfer of performance to the competitive situation been measured? Correlational and anecdotal accounts should be interpreted with caution.

The Sport and Exercise Scientist • Issue 38 • Winter 2013 • www.bases.org.uk



Nicholas is a Principal Lecturer in Sport and Exercise Psychology at the University of Brighton.



## Dr Jenny Page

lenny is a Senior Lecturer in Sport and Exercise Psychology at the University of Chichester

Dr Mark Wilson

Movement Science at



Dr Joe Causer

oe is a Senior Lecturer n Applied Sport Psychology at Liverpool ohn Moores University.



Dr Rob Gray

Rob is a Reader in Perception and Action at Birmingham University.



the University of Exeter.

Mark is Professor and subject leader of sport sciences at Brunel University

The authors would like to acknowledge the Expertise and Skill Acquisition Network (ESAN) for helping to bring this team of researchers together.

Abernethy, B. (1996). Training the visual-perceptual skills of athletes - Insights from the study of motor expertise. American Journal of Sports Medicine, 24, S89-S92.

Abernethy, B. & Wood, J.M. (2001). Do generalized visual training programmes for sport really work? An experimental investigation. Journal of Sports Sciences, 19(3), 203-222. doi: 10.1080/026404101750095376

Causer, J. et al. (2012). Perceptual training: What can be trained? In N.J. Hodges & A.M. Williams (Eds.), Skill Acquisition in Sport: Research, Theory and Practice (pp. 306-324), London: Routledge.

Clark, J.F. et al. (2012). High-performance vision training improves batting statistics for University of Cincinnati baseball players. Plos One, 7(1). doi:10.1371/journal.

Goodrich, G.L. et al. (2013). Development of a mild traumatic brain injury-specific vision screening protocol: A Delphi study. Journal of Rehabilitation Research and Development 50(6) 757-768 doi: 10 1682/irrd 2012 10 0184

Hopwood, M.J. et al. (2011). Does visual-perceptual training augment the fielding performance of skill cricketers? International Journal of Sports Science & Coaching, 4(6),

Schwab, S. & Memmert, D. (2012). The impact of a sports vision training program in youth field hockey players. Journal of Sports Science and Medicine, 11(4), 624-631.

Smeeton, N.J. et al. (2005). The relative effectiveness of various instructional approaches in developing anticipation skill. Journal of Experimental Psychology-Applied, 11(2), 98-110. doi: 10.1037/1076-898x.11.2.98.

Vine, S.J., Moore, L.J. & Wilson, M.R. Quiet eye training: The acquisition, refinement and resilient performance of targeting skills. European Journal of Sport Science ahead-of-print (2012): 1-8. doi:10.1080/17461391.2012.683815

Williams, A.M. & Grant, A. (1999). Training perceptual skill in sport. International Journal of Sport Psychology, 30(2), 194-220.

Williams, A.M. et al. (2002). Anticipation skill in a real-world task: Measurement, training, and transfer in tennis. Journal of Experimental Psychology-Applied, 8(4), 259-270. doi:10.1037/1076-898x.8.4.259

Download a PDF of this article: www.bases.org.uk/BASES-Expert-Statements

#### Copyright © BASES, 2013

mission is given for reproduction in substantial part. We ask that the following note be included: "First published in The Sport and Exercise Scientist, Issue 38, Winter 2013. Published by the British Association of Sport and Exercise Sciences – www.bases.org.uk"