

In press at Alcoholism and Drug Addiction.

Spatial Anticipatory Attentional Bias for Alcohol: A Preliminary Report on Reliability and Associations with Risky Drinking

Thomas E. Gladwin ^{a*}

^a Department of Psychology & Counselling, University of Chichester. College Lane, Chichester, PO19 6PE, United Kingdom. Email: thomas.gladwin@gmail.com. Tel.: +44(0)7895625183.

* Corresponding author.

Author notes

Declarations of interest: none.

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

The manuscript has been available as a preprint on PsyArXiv.

Abstract

Introduction

Although risky drinking and alcohol dependence have been associated with spatial attentional biases, concerns have been raised about the reliability of the frequently-used dot-probe task. A form of anticipatory bias related to predictive cues has been found to be related to alcohol-related processes, and to have high reliability in the context of threat stimuli. It remains to be determined whether this anticipatory attentional bias also has good reliability for alcohol stimuli. Further, correlations with drinking-related individual differences need to be replicated.

Material and Methods

83 healthy adult participants were included, who completed the task and questionnaires on risky drinking (AUDIT-C), drinking motives (DMQ-R), reasons to abstain from drinking (RALD), and alcohol craving (ACQ). The task used a 400 ms Cue-Stimulus Interval, based on previous work. The Spearman-Brown split-half reliability of reaction time-based bias scores was calculated. The within-subject effect of probe location (predicted-alcohol versus predicted-non-alcohol) was tested using a paired-sample t-test. Correlations were calculated between bias scores and questionnaire scales; tests were one-sided for predicted effects and two-sided for exploratory effects.

Results

The reliability was .81 (.74 after outlier removal). There was no overall bias. A predicted correlation between risky drinking and anticipatory bias towards alcohol was found, but no other predicted or exploratory effects.

Discussion

The anticipatory attentional bias for alcohol is a reliably measurable individual difference, with some evidence that it is associated with risky drinking.

Conclusions

Implicit behavioural measures of spatial attentional bias can in principle achieve high reliability. Further study of attentional biases using predictive cues would appear to be promising.

Keywords

Dot-probe task; attentional bias; anticipatory attentional bias; alcohol; reliability; risky alcohol use

Acknowledgments

There was no funding for this study.

Introduction

The dot-probe task [1,2] is often used to measure spatial attentional biases. In trials of this task, first two cues are shown, from two different categories such as alcohol versus soft drink, and subsequently probe stimuli are presented at the location of one of the cues. Differences in responses to the probes that depend on the category of the cues at their location suggest that the cues caused a bias in processing; e.g., if one of the cue types tends to capture attention, then responses to probes at its location should be faster. Alcohol-related stimulus have been found to affect spatial attention [3–6]. In individuals reporting heavy social drinking attention appears to be drawn towards alcohol cues [4,5], as would be expected given theory on incentive salience [7,8]. However, complex, time-dependent patterns have been found in dependence, with a shift from initial orienting to attentional disengagement [6,9,10].

Further, a problem with the dot-probe task is that it has been found to have low reliability in a number of studies [11–19]. This would seem to pose a serious threat to at least some forms of attentional bias research using behavioural measures. If the current literature is taken to imply that any behavioural measure of attentional bias is unacceptably likely to be noisy, this could lead to a shift to interesting but expensive and less widely accessible psychophysiological methods. It is essential to ask whether this shift is truly necessary and whether it is justified to consider behavioural measures inherently problematic [20].

One alternative approach is to explore novel versions of the dot-probe task: perhaps a reliable behavioural measure is possible but not yet known. One such variant of the task was suggested by the R3 model – the Reprocessing and Reinforcement model of Reflectivity [21,22]. In this model, automatic attentional biases can occur due to the predicted outcomes of attentional shifts (note that “prediction” here is conceived of as a low-level, underlying process caused by prior reinforcement learning; rather than an effortful attempt to make a prediction about the future). For example, if an individual believes that something scary is lurking behind a door, the anticipation of what could happen on opening the door will affect their attention towards it. Or, if there might be

something tasty out of sight in a cupboard, the predicted outcome of finding a treat might involuntarily affect attention. It thus seems that attentional biases could occur in response to information on what will likely happen if attention is directed to one location or another: will something attractive or aversive appear at that location? And could that kind of prediction of outcome automatically affect attentional shifts? A cued version of the dot-probe task, termed the cued Visual Probe Task, cVPT [23], was developed to measure such anticipatory or outcome-related effects. In this task, instead of presenting emotional cues intended to evoke an automatic stimulus-driven response, two visually neutral predictive cues are presented at the start of each trial. On one half of the trials, the predictive cues are replaced by affective stimuli, one cue always being replaced by a stimulus from one category (e.g., alcohol or threat) and the other by a stimulus from another category (e.g., water or safe). On the other half of the trials, probe stimuli are presented requiring a response; on these trials, no affective stimuli are presented at all. Thus, performance on probes is never influenced by the direct presentation of an emotional stimulus, only by the location of visually neutral cues predicting stimulus categories. Further, it has been found that performing a training version of the cVPT induces an attentional bias to stimuli belonging to the trained predicted categories [24]. This supports the interpretation of effects on the cVPT being due to anticipatory processes. It could be expected that performing the cVPT only involves the visually neutral cues acquiring salience, rather than outcome-related processes. However, this would not be expected to lead to the predicted stimulus categories acquiring a bias following the cVPT-training, rather than just the predictive cues.

In the context of threat-related attentional bias, reliability was found to be improved in the cVPT relative to a usual VPT in which emotional cues were presented before probes [25]. The improved reliability was suggested to be due to the removal or mitigation of noisy influences that could play a role when actually presenting emotional stimuli, such as varying responses to particular exemplars, or potentially complex patterns of multiple cognitive responses to actually presented emotional stimuli. While the anticipatory attentional bias has been previously used to study alcohol-related

attentional bias and bias variability [23,26], its reliability has not yet been evaluated for alcohol-related stimuli. Further, the validity of the bias as a reflection of processes related to alcohol and risky drinking requires additional support.

The current study therefore aimed, first, to determine the reliability of the anticipatory attentional bias for alcohol; and, second, to explore correlations between the anticipatory bias and alcohol-related individual differences.

Material and method

Participants

The experiment was completed online by an analytical sample of 83 healthy adult participants (77 female, 9 male; age 19.7, $SD = 2.95$). A further 15 participants performed the experiment but were excluded due to inadequate performance (mean accuracy below .90) indicating, given the simplicity of the tasks and usual accuracy levels, that these participants were not performing the task as required. Participants gave informed consent, and the study was approved by the institutional ethics committee.

Questionnaires

The following questionnaires were used to assess individual differences related to alcohol use. The 3-item Alcohol Use Disorders Identification Test - Consumption, AUDIT-C, was used to measure hazardous drinking [27–30]. The Drinking Motives Questionnaire Revised, DMQ-R [31] provided measures of Enhancement, Social, Coping, and Conformity motives. The Reasons for Abstaining or Limiting Drinking questionnaire, RALD [32,33] was used to measure motives to refrain from drinking: Loss of Control, Adverse Consequences, and Convictions. Finally, aspects of craving for alcohol were measured with the Alcohol Craving Questionnaire – Short Form, ACQ [34,35]: Compulsivity, Expectancy, Purposefulness and Emotionality.

Tasks

The cued Visual Probe Task consisted of 9 blocks of 40 trials, preceded by a training phase of 8 blocks of 40 trials. Trials were divided into Picture and Probe trials.

Probe trials started with the presentation of a central fixation cross for 300, 400 or 500 ms, followed by the presentation of two visually neutral cues: the symbols O O O O and | | | |, presented in yellow versus blue. The cues were onscreen for a Cue-Stimulus Interval of 400 ms. The cues were located on alternating diagonals per trial: either on the top-left and bottom-right of the screen, or on the bottom-left and top-right of the screen. Following the cue period, a probe stimulus, >><<, was presented at one of the cue locations, and a distractor stimulus, /\ or \/, at the other location. The probe stimulus was presented for 1000 ms, or until a response was given if faster than 1000 ms. The task was to quickly and accurately press a key (R, F, I or J) corresponding to the probe location. Errors were followed by a red -1 for incorrect responses, and a red "Too late!" if no response was given, for 200 ms.

On Picture trials, the cues were replaced by pictures presented at the cue locations. One of the cues was always replaced by an alcoholic stimulus (a color picture of a glass or bottle of an alcoholic beverage), centered on the cue location. The other cue was always replaced by a non-alcoholic stimulus (a color picture of a non-alcoholic beverage). The mapping of cues to stimulus category was randomized over subjects. The pictures remained onscreen for 1000 ms, followed by 200 ms of empty screen.

Procedure

Participants performed the experiment fully online. They first filled in the questionnaires and subsequently completed the training and assessment phase of the task. For transparency, we note that participants performed additional tasks and sessions unrelated to the current study.

Preprocessing and statistical analyses

Trials were removed that were likely not to reflect normal task performance: the first four trials of the task, trials following an error, and the first trial of each block. Of the remaining probe trials, the median reaction time was calculated for the Probe-on-Alcohol and Probe-on-Non-alcohol predictive cue locations. The reliability of the bias, i.e., the median RT for the Alcohol minus Non-alcohol locations, was tested using Spearman-Brown formula for the split-half reliability of the task, which

was divided in sets of even and odd numbered blocks. The effect of probe location was tested using paired-sample t-tests, and correlations were tested between questionnaire scores and the individual bias scores. Based on previous results involving the same Cue-Stimulus Interval of 400 ms [26], we could hypothesize an increasing bias towards alcohol (i.e., more negative bias scores) with increasing scores on the AUDIT, ACQ-Compulsivity, ACQ-Expectancy, and RALD-Convictions and therefore used one-sided tests ($p < .1$ criterion) for these scales.

Data and software will be made available upon request.

Results

The mean sample scores on the questionnaires are given in Table 1. Using cut-offs for risky drinking in a student population [36], the percentage of female participants with an AUDIT-C score of at least 5 was 35%, and the percentage of male participants with an AUDIT-C score of at least 7 was 25%.

Table 1. Questionnaire scores

Scale	Mean (SD)
AUDIT-C	4.30 (2.23)
DMQ-Social	2.94 (0.97)
DMQ_Coping	1.70 (0.79)
DMQ-Enhancement	2.45 (1.03)
DMQ-Conformity	1.52 (0.71)
ACQ-Compulsivity	1.20 (0.58)
ACQ-Expectancy	1.97 (1.04)
ACQ-Purposefulness	4.57 (1.49)
ACQ-Emotionality	1.85 (1.09)

The reliability of bias scores was .84. Removal of extreme points (z -score > 3 on either the “even” or “odd” bias, $n = 2$) resulted in a reliability of .74.

There was no overall effect of probe location, with reaction times of 486 ms ($SD = 48$ ms) for probes on the non-alcohol location and 489 ms ($SD = 44$ ms) for probes on the alcohol location. Of the a priori expected one-sided relationships with questionnaires, only higher AUDIT scores were associated with a bias towards alcohol, $r = -0.22$, $p = 0.050$. There were no other significant correlations.

Discussion

The current results agree with previous studies in two main ways. First, the anticipatory attentional bias for alcohol was found to have good split-half reliability. This is of interest in relation to the concern with reliability for the usual dot-probe task [18]. The current task's reliance on only anticipatory attentional processes may aid reliability: any interfering processes evoked by actual stimulus presentation do not occur on probe trials, and there is no variability due to the immediate presentation of different specific exemplars (although there could of course still be more complex history effects related to the particular sequence of presented stimuli). Any bias must be due to the learned relationship between the simple, non-varying cues and the overall stimulus categories. The diagonalized form of the task may also contribute as neither responses nor stimulus locations were ever immediately repeated, reducing potential trial-to-trial carryover effects.

The availability of a form of spatial attentional bias with good reliability would be of interest when studying correlations between attentional bias and other individual differences, or when the bias is used as a dependent variable in a training study. Further, this finding indicates that the anticipatory bias is a consistent individual difference: individuals systematically differ in the degree to which their attention is affected by predicted outcomes of attentional shifts towards or away from cued locations. The current findings may thus be useful for research aimed at a better understanding of the nature of attentional biases and automaticity, and of the influences of task variations on psychometric properties.

Second, supporting the interpretation of individual differences in the bias in terms of attention to predicted outcome categories and hence the validity of the bias, hazardous drinking was indeed associated with an anticipatory bias towards predicted alcohol location, as in a previous study [26]. However, as other predicted effects were not found, and the effect size of the current result was not large and the test would not survive correction for multiple testing, future research should focus on replicating the specific correlation between risky drinking and anticipatory attentional bias. The relatively weak correlation between bias and risky drinking may have been due to the limitation of a low overall level of risky drinking in the current sample. Further research is now necessary in larger samples with a wider range of risky drinking, and in clinical groups as compared to control groups. Another important limitation of the current opportunistic sample is the unequal distribution of male and female participants. Future work should consider moderating effects of gender on the relationships between anticipatory bias and alcohol-related individual differences.

Another limitation is the use of only a single Cue-Stimulus Interval, of 400 ms. While this was selected based on previous work, it must be acknowledged that the current results are specific to this interval, and longer or shorter durations could well produce different reliabilities and effects due to the role of temporal dynamics in attentional biases [6,9,10].

Conclusions

The current results provide further support for the concept and measurability of an anticipatory attentional bias for alcohol. Risky drinking may be related to enhanced salience of predicted alcohol stimuli. More generally, biases can be due to the selection of cognitive responses, such as attentional shifts, based on their predicted outcome; e.g., whether they are likely to focus attention on upcoming salient stimuli. The results also add to the evidence suggesting that behavioural measures of attentional bias can in principle be both reliable and valid, but it may be necessary to use novel variations of tasks and concepts.

References

1. Mogg K, Bradley BP. Orienting of Attention to Threatening Facial Expressions Presented under Conditions of Restricted Awareness. *Cogn Emot*, 1999;13(6):713–40.
2. MacLeod CM, Mathews A, Tata P. Attentional bias in emotional disorders. *J Abnorm Psychol*, 1986;95(1):15–20.
3. Field M, Cox WM. Attentional bias in addictive behaviors: a review of its development, causes, and consequences. *Drug Alcohol Depend*, 2008;97(1–2):1–20.
4. Field M, Mogg K, Zetteler J, Bradley BP. Attentional biases for alcohol cues in heavy and light social drinkers: the roles of initial orienting and maintained attention. *Psychopharmacology (Berl)*, 2004;176(1):88–93.
5. Townshend JM, Duka T. Attentional bias associated with alcohol cues: differences between heavy and occasional social drinkers. *Psychopharmacology (Berl)*, 2001;157(1):67–74.
6. Townshend JM, Duka T. Avoidance of alcohol-related stimuli in alcohol-dependent inpatients. *Alcohol Clin Exp Res*, 2007;31(8):1349–57.
7. Volkow ND, Koob GF, McLellan AT. Neurobiologic Advances from the Brain Disease Model of Addiction. *N Engl J Med*, 2016;374(4):363–71.
8. Robinson TE, Berridge KC. Addiction. *Annu Rev Psychol*, 2003;54:25–53.
9. Noël X, Colmant M, Van Der Linden M, Bechara A, Bullens Q, Hanak C, et al. Time course of attention for alcohol cues in abstinent alcoholic patients: the role of initial orienting. *Alcohol Clin Exp Res*, 2006;30(11):1871–7.
10. Vollstädt-Klein S, Loeber S, von der Goltz C, Mann K, Kiefer F. Avoidance of alcohol-related stimuli increases during the early stage of abstinence in alcohol-dependent patients. *Alcohol*, 2009;44(5):458–63.

11. Brown HM, Eley TC, Broeren S, MacLeod CM, Rinck M, Hadwin JA, et al. Psychometric properties of reaction time based experimental paradigms measuring anxiety-related information-processing biases in children. *J Anxiety Disord*, 2014;28(1):97–107.
12. Dear BF, Sharpe L, Nicholas MK, Refshauge K. The psychometric properties of the dot-probe paradigm when used in pain-related attentional bias research. *J Pain*, 2011;12(12):1247–54.
13. Puls S, Rothermund K. Attending to emotional expressions: no evidence for automatic capture in the dot-probe task. *Cogn Emot*, 2017;1–14.
14. Schmukle SC. Unreliability of the dot probe task. *Eur J Pers*, 2005;19(7):595–605.
15. Waechter S, Nelson AL, Wright C, Hyatt A, Oakman J. Measuring attentional bias to threat: Reliability of dot probe and eye movement indices. *Cognit Ther Res*, 2014;38(3):313–33.
16. Kappenman ES, Farrens JL, Luck SJ, Proudfit GH. Behavioral and ERP measures of attentional bias to threat in the dot-probe task: Poor reliability and lack of correlation with anxiety. *Front Psychol*, 2014;5(DEC):1368.
17. McNally RJ. Attentional bias for threat: Crisis or opportunity? *Clin Psychol Rev*, 2018;
18. Christiansen P, Schoenmakers TM, Field M. Less than meets the eye: Reappraising the clinical relevance of attentional bias in addiction. *Addict Behav*, 2015;44:43–50.
19. Ataya AF, Adams S, Mullings E, Cooper RM, Attwood AS, Munafò MR. Internal reliability of measures of substance-related cognitive bias. *Drug Alcohol Depend*, 2012;121(1–2):148–51.
20. MacLeod CM, Grafton B, Notebaert L. Anxiety-Linked Attentional Bias: Is It Reliable? *Annu Rev Clin Psychol*, 2019;15.
21. Gladwin TE, Figner B, Crone EA, Wiers RW. Addiction, adolescence, and the integration of control and motivation. *Dev Cogn Neurosci*, 2011;1(4):364–76.
22. Gladwin TE, Figner B. “Hot” cognition and dual systems: Introduction, criticisms, and ways

- forward. In: Wilhelms E, Reyna VF, editors. *Frontiers of Cognitive Psychology Series: Neuroeconomics, Judgment and Decision Making*, New York: Psychology Press; 2014. p. 157–80.
23. Gladwin TE. Attentional bias variability and cued attentional bias for alcohol stimuli. *Addict Res Theory*, 2016;25(1):32–8.
 24. Gladwin TE, Möbius M, Becker ES. Predictive Attentional Bias Modification induces stimulus-evoked attentional bias for threat. *Eur J Psychol*, 2019;In press.
 25. Gladwin TE, Möbius M, Mcloughlin S, Tyndall I. Anticipatory versus reactive spatial attentional bias to threat. *Br J Psychol*, 2019;110(1):3–14.
 26. Gladwin TE, Vink M. Alcohol-related attentional bias variability and conflicting automatic associations. *J Exp Psychopathol*, 2018;9(2).
 27. Bush K, Kivlahan DR, McDonnell MB, Fihn SD, Bradley KA. The AUDIT alcohol consumption questions (AUDIT-C): an effective brief screening test for problem drinking. Ambulatory Care Quality Improvement Project (ACQUIP). Alcohol Use Disorders Identification Test. *Arch Intern Med*, 1998;158(16):1789–95.
 28. Bradley KA, DeBenedetti AF, Volk RJ, Williams EC, Frank D, Kivlahan DR. AUDIT-C as a brief screen for alcohol misuse in primary care. *Alcohol Clin Exp Res*, 2007;31(7):1208–17.
 29. Gordon AJ, Maisto SA, McNeil M, Kraemer KL, Conigliaro RL, Kelley ME, et al. Three questions can detect hazardous drinkers. *J Fam Pract*, 2001;50(4):313–20.
 30. Gual A, Segura L, Contel M, Heather N, Colom J. Audit-3 and audit-4: effectiveness of two short forms of the alcohol use disorders identification test. *Alcohol Alcohol*, 2002;37(6):591–6.
 31. Cooper ML. Motivations for alcohol use among adolescents: Development and validation of a

- four-factor model. *Psychol Assess*, 1994;6(2):117–28.
32. Epler AJ, Sher KJ, Piasecki TM. Reasons for abstaining or limiting drinking: a developmental perspective. *Psychol Addict Behav*, 2009;23(3):428–42.
 33. Anderson KG, Grunwald I, Bekman N, Brown SA, Grant A. To drink or not to drink: Motives and expectancies for use and nonuse in adolescence. *Addict Behav*, 2011;36(10):972–9.
 34. Singleton E, Henningfield J, Heishman T, Douglas E, Tiffany S. Multidimensional aspects of craving for alcohol. In: Proceedings of the 57th Annual Meeting of the College on Problems of Drug Dependence, 1995.
 35. Connolly KM, Coffey SF, Baschnagel JS, Drobles DJ, Saladin ME. Evaluation of the Alcohol Craving Questionnaire-Now factor structures: application of a cue reactivity paradigm. *Drug Alcohol Depend*, 2009;103(1–2):84–91.
 36. DeMartini KS, Carey KB. Optimizing the use of the AUDIT for alcohol screening in college students. *Psychol Assess*, 2012;24(4):954–63.